

Patient Name _____

Examiner Name _____ Date/Time of Exam _____



STANDARD NEUROLOGICAL CLASSIFICATION OF SPINAL CORD INJURY



MOTOR

KEY MUSCLES
(scoring on reverse side)

| | | | |
|----|--------------------------|--------------------------|--|
| | R | L | |
| C5 | <input type="checkbox"/> | <input type="checkbox"/> | Elbow flexors |
| C6 | <input type="checkbox"/> | <input type="checkbox"/> | Wrist extensors |
| C7 | <input type="checkbox"/> | <input type="checkbox"/> | Elbow extensors |
| C8 | <input type="checkbox"/> | <input type="checkbox"/> | Finger flexors (distal phalanx of middle finger) |
| T1 | <input type="checkbox"/> | <input type="checkbox"/> | Finger abductors (little finger) |

UPPER LIMB TOTAL (MAXIMUM) + =
(25) (25) (50)

Comments:

| | | | |
|----|--------------------------|--------------------------|-----------------------|
| L2 | <input type="checkbox"/> | <input type="checkbox"/> | Hip flexors |
| L3 | <input type="checkbox"/> | <input type="checkbox"/> | Knee extensors |
| L4 | <input type="checkbox"/> | <input type="checkbox"/> | Ankle dorsiflexors |
| L5 | <input type="checkbox"/> | <input type="checkbox"/> | Long toe extensors |
| S1 | <input type="checkbox"/> | <input type="checkbox"/> | Ankle plantar flexors |

Voluntary anal contraction (Yes/No)

LOWER LIMB TOTAL (MAXIMUM) + =
(25) (25) (50)

| | LIGHT TOUCH | | PIN PRICK | |
|------|-------------|---|-----------|---|
| | R | L | R | L |
| C2 | | | | |
| C3 | | | | |
| C4 | | | | |
| C5 | | | | |
| C6 | | | | |
| C7 | | | | |
| C8 | | | | |
| T1 | | | | |
| T2 | | | | |
| T3 | | | | |
| T4 | | | | |
| T5 | | | | |
| T6 | | | | |
| T7 | | | | |
| T8 | | | | |
| T9 | | | | |
| T10 | | | | |
| T11 | | | | |
| T12 | | | | |
| L1 | | | | |
| L2 | | | | |
| L3 | | | | |
| L4 | | | | |
| L5 | | | | |
| S1 | | | | |
| S2 | | | | |
| S3 | | | | |
| S4-5 | | | | |



TOTALS { + = } + =
(MAXIMUM) (56) (56) (56) (56)

Any anal sensation (Yes/No)
 PIN PRICK SCORE (max: 112)
 LIGHT TOUCH SCORE (max: 112)

SENSORY

KEY SENSORY POINTS

0 = absent
1 = impaired
2 = normal
NT = not testable



• Key Sensory Points

| | | | | | | | | | |
|---|---------|--------------------------|--------------------------|---|--------------------------|---|---------|--------------------------|--------------------------|
| NEUROLOGICAL LEVEL <small>The most caudal segment with normal function</small> | SENSORY | R | L | COMPLETE OR INCOMPLETE? <small>Incomplete = Any sensory or motor function in S4-S5</small> | <input type="checkbox"/> | ZONE OF PARTIAL PRESERVATION <small>Caudal extent of partially innervated segments</small> | SENSORY | R | L |
| | MOTOR | <input type="checkbox"/> | <input type="checkbox"/> | | | | MOTOR | <input type="checkbox"/> | <input type="checkbox"/> |
| ASIA IMPAIRMENT SCALE | | | | <input type="checkbox"/> | | | | | |

MUSCLE GRADING

- 0 total paralysis
- 1 palpable or visible contraction
- 2 active movement, full range of motion, gravity eliminated
- 3 active movement, full range of motion, against gravity
- 4 active movement, full range of motion, against gravity and provides some resistance
- 5 active movement, full range of motion, against gravity and provides normal resistance
- 5* muscle able to exert, in examiner's judgement, sufficient resistance to be considered normal if identifiable inhibiting factors were not present

NT not testable. Patient unable to reliably exert effort or muscle unavailable for testing due to factors such as immobilization, pain on effort or contracture.

ASIA IMPAIRMENT SCALE

- A = Complete:** No motor or sensory function is preserved in the sacral segments S4-S5.
- B = Incomplete:** Sensory but not motor function is preserved below the neurological level and includes the sacral segments S4-S5.
- C = Incomplete:** Motor function is preserved below the neurological level, and more than half of key muscles below the neurological level have a muscle grade less than 3.
- D = Incomplete:** Motor function is preserved below the neurological level, and at least half of key muscles below the neurological level have a muscle grade of 3 or more.
- E = Normal:** Motor and sensory function are normal.

CLINICAL SYNDROMES (OPTIONAL)

- Central Cord
- Brown-Sequard
- Anterior Cord
- Conus Medullaris
- Cauda Equina

STEPS IN CLASSIFICATION

The following order is recommended in determining the classification of individuals with SCI.

1. Determine sensory levels for right and left sides.
2. Determine motor levels for right and left sides.
Note: in regions where there is no myotome to test, the motor level is presumed to be the same as the sensory level.
3. Determine the single neurological level.
This is the lowest segment where motor and sensory function is normal on both sides, and is the most cephalad of the sensory and motor levels determined in steps 1 and 2.
4. Determine whether the injury is Complete or Incomplete (sacral sparing).
If voluntary anal contraction = No AND all S4-5 sensory scores = 0 AND any anal sensation = No, then injury is COMPLETE. Otherwise injury is incomplete.

5. Determine ASIA Impairment Scale (AIS) Grade:

Is injury Complete?

NO



Is injury motor incomplete?

YES



Are at least half of the key muscles below the (single) neurological level graded 3 or better?

NO



AIS=C

YES



AIS=D

If sensation and motor function is normal in all segments, AIS=E
Note: AIS E is used in follow up testing when an individual with a documented SCI has recovered normal function. If at initial testing no deficits are found, the individual is neurologically intact; the ASIA Impairment Scale does not apply.